

The National Immunisation Schedule



| Age | Disease to protect against | Vaccine |
|-------------|--|--|
| Pregnancy | Tetanus + diphtheria + whooping cough (pertussis) | Boostrix® (28 to 38 weeks) |
| | Influenza | Any time during pregnancy. |
| 6 Weeks | Rotavirus (first dose must be given before 15 weeks) | Rotarix® (oral) |
| | Diphtheria + tetanus + whooping cough (pertussis) + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib) | Infanrix® hexa |
| | Pneumococcal disease | Synflorix® |
| 3 Months | Rotavirus (second dose must be given before 25 weeks) | Rotarix® (oral) |
| | Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib) | Infanrix® hexa |
| | Pneumococcal disease | Synflorix® |
| 5 Months | Diphtheria + tetanus + whooping cough + polio + hepatitis B + <i>Haemophilus influenzae</i> type b (Hib) | Infanrix® hexa |
| | Pneumococcal disease | Synflorix® |
| 15 Months | <i>Haemophilus influenzae</i> type b (Hib) | Hiberix® |
| | Measles + mumps + rubella | Priorix® |
| | Pneumococcal disease | Synflorix® |
| | Chickenpox (varicella) | Varilrix® |
| 4 Years | Diphtheria + tetanus + whooping cough + polio | Infanrix® IPV |
| | Measles + mumps + rubella | Priorix® |
| 11+12 Years | Tetanus + diphtheria + whooping cough | Boostrix® |
| | Human papillomavirus (HPV) | Gardasil®9 (2 doses, 6 months apart) |
| 45 Years | Tetanus + diphtheria | ADT® Booster |
| 65 Years | Tetanus + diphtheria | ADT® Booster |
| | Influenza | Given annually. |
| | Shingles | Zostavax® |